


51 CHEMISTRY REACTIONS

You Can't Afford to MISS!


1. Aldol condensation

Two molecules of aldehydes or ketones having α -hydrogen condense together in presence of base [dil. NaOH, Na_2CO_3 , $\text{Ba}(\text{OH})_2$] to form β -hydroxyaldehyde or β -hydroxyketone respectively which are collectively known as **aldol**.




2. Birch Reduction

In case of disubstituted alkynes reduction with sodium in liq. ammonia gives *trans*-alkene predominantly. This type of reduction is known as Birch-reduction.




3. Beckmann Rearrangement

Oximes on treatment with catalyst such as conc. H_2SO_4 , SOCl_2 , PCl_5 , etc. undergo rearrangement to form substituted amides. Reaction is known as 'Beckmann rearrangement'.





4. Balz-Schiemann's reaction

This is the method to prepare fluorobenzene from benzene diazonium salt by first converting it to benzene diazonium tetrafluoroborate.




5. Clemmensen Reduction

In this method, aldehydes and ketones are reduced with Zn - Hg amalgam and conc. HCl.




6. Corey-House synthesis

In this reaction, an alkyl halide (1° , 2° or 3°) is first converted into lithium dialkyl copper and then reacted with another alkyl halide (preferably 1°) to give corresponding alkane.




7. Cope elimination

Tertiary amines are not oxidised by KMnO_4 . Their oxidation is carried out with Caro's acid or ozone or hydrogen peroxide to form amine oxide.




Tertiary amine oxide containing a β -hydrogen atom forms an alkene. This reaction is called **Cope reaction** or **Cope elimination**.




8. Carbylamine Reaction

Primary amines both aliphatic and aromatic on heating with chloroform and alcoholic solution of potassium hydroxide form isocyanides or carbylamines which have an extremely unpleasant odour.




9. Claisen—Schmidt reactions

It is a base catalyzed crossed aldol condensation in which an aromatic aldehyde condenses with aliphatic aldehyde or ketone. It is also known as simply **Claisen reaction**.




10. Cannizzaro Reaction

It is a self oxidation-reduction reaction in which aldehydes that do not have any α -hydrogen atom undergo disproportionation reaction (*i.e.* self-redox reaction) in the presence of 50% aqueous or ethanolic solution of alkali in which one of the molecule being reduced to alcohol and other being oxidised to the salt of corresponding acid.




11. Claisen condensation

Esters containing α -hydrogen atoms undergo self-condensation in presence of strong base like sodium ethoxide to give β -keto esters.




12. Curtius rearrangement

In this method an acyl azide (obtained from reaction of carboxylic acid chloride and sodium azide), on heating in acidic or alkaline medium yields primary amine. The reaction involves an intermediate isocyanate which is hydrolysed to give primary amine.




13. Diels-Alder Reaction

This reaction involves the addition of a conjugated diene (4π -electron system) to an unsaturated compound called dienophile (2π -electron system) to yield a cyclic system.




14. Etard Reaction

Aromatic hydrocarbons on treatment with mild oxidising agents such as chromyl chloride CrO_2Cl_2 yield aldehydes.



15. Frankland Reaction

This is similar to Wurtz reaction with the difference that instead of sodium, zinc is used here in inert solvent.




Dialkyl zinc compounds react with tertiary alkyl halides to give hydrocarbons with quaternary carbon atoms only.




16. Finkelstein reaction

Iodoalkanes can be conveniently prepared by treating alkyl chloride or bromides with sodium iodide in acetone or methanol. This is known as Finkelstein reaction.




17. Friedel-Crafts Reaction

Simple aromatic hydrocarbons can be converted to substituted hydrocarbons by reaction with alkyl halide in presence of Lewis acid like AlCl_3 or FeCl_3 , etc.



18. Fries Rearrangement


Phenyl esters on heating with anhydrous AlCl_3 undergo Fries rearrangement in which acyl group (acetyl, benzoyl, etc.) migrates from phenolic oxygen to *ortho* or *para* position of benzene ring resulting in the formation of mixture of *o*- and *p*-hydroxyketone.



19. Gabriel phthalimide synthesis


Phthalimide on reaction with caustic potash forms potassium phthalimide which on treatment with alkyl halide gives *N*-alkyl phthalimide. This *N*-alkyl phthalimide on hydrolysis with hydrochloric acid yields primary amines.






30 Meerwein-Ponndorf-Verley reduction

In this reaction, ketones are reduced to secondary alcohols on heating with *iso*-propyl alcohol in presence of catalyst aluminium isopropoxide. In the reaction *iso*-propyl alcohol gets oxidised to acetone.




31. Ozonolysis

Ozone on passing through a solution of alkenes in an inert solvent like CCl₄ or CHCl₃ at low temperature oxidises the alkene to a cyclic product called ozonide which is unstable and on reduction with reducing agents like zinc and water or hydrogen in presence of palladium gives aldehydes and/or ketone depending upon structure of alkene, i.e., >C= part of the alkene gets converted to >C=O.




32. Oppenauer's oxidation

In this reaction, secondary alcohols are oxidised to ketones in a good yield using aluminium *tert*-butoxide in excess of acetone.




33. Oxo process

In this process, an alkene on heating with water gas (CO + H₂) under pressure in the presence of catalyst cobalt carbonyl hydride [CoH(CO)₄] yields aldehyde.




34. Perkin condensation

It is a condensation reaction in which an aromatic aldehyde condenses with aliphatic acid anhydride in presence of sodium salt of same acid to give a condensate which upon hydrolysis yields α,β-unsaturated acid.




35. Reimer-Tiemann reaction

Concentrated solution of CHCl₃ reacts with phenol in presence of alc. KOH to yield salicylaldehyde.




36. Reformatsky reaction

It involves the reaction of zinc and α-halogenated ester, which add to carbonyl compound to yield β-hydroxy ester.




37. Rosenmund's reduction

Acid chlorides are reduced to aldehydes by palladised barium sulphate partially poisoned with sulphur or quinoline. (–COCl is reduced to –CHO)




38. Sandmeyer's reaction

In this reaction Diazonium salt reacts with CuX/HX and respective halide is formed.




39. Schotten-Baumann reaction



Benzoylation of alcohols or phenols in presence of NaOH is called **Schotten-Baumann reaction**.


40. Stephen's reduction

In this reaction cyanides are reduced to aldehyde in pressure of SnCl_2/H_4 followed by hydrolysis.




41. Schmidt's reaction

Carboxylic acids on heating with hydrazoic acid (HN_3) at $50-55^\circ\text{C}$ in presence of conc. H_2SO_4 produce primary amines with one carbon atom less than original carboxylic acid.




42. Sabatier - Senderen's reaction

The method involves the reduction of unsaturated hydrocarbons (alkenes and alkynes) with nickel catalyst. Raney nickel is more reactive than supported nickel catalyst.




43. Tischenko reaction

All aldehydes (with or without α -hydrogen atom) in presence of aluminium ethoxide, $[\text{Al}(\text{OC}_2\text{H}_5)_3]$ undergo simultaneous oxidation (to carboxylic acid) and reduction (to alcohol) to form an ester.




44. Ullmann Reaction

Aryl halides especially iodides on reaction with copper metal yield biphenyls.




45. Williamson's synthesis

Metal alkoxides on treatment with alkyl halides yield ethers. The reaction is known as Williamson's synthesis.




46. Wacker process

Alkenes are directly oxidised by the catalyst palladium chloride-cupric chloride ($\text{PdCl}_2\text{-CuCl}_2$) in the presence of air or oxygen.




47. Wolff-Kishner Reduction

In this method, aldehydes and ketones are reduced with hydrazine ($\text{NH}_2\text{-NH}_2$) and KOH.



48. Wurtz Reaction

In this reaction, ether solution of alkyl halide is treated with sodium to form alkane.




49. Wittig Reaction

It is an important reaction to change a carbonyl compound into alkene by treating carbonyl compound with a ylide. Ylides are the species which in their ground state have negative carbon adjacent to a positive heteroatom like P, As, Sb, etc. For example, phosphorus ylide, $\text{Ph}_3\text{P}^+ - \text{CHR}^-$. $\text{R}'\text{CH}=\text{O} + \text{Ph}_3\text{P}^+ - \text{CHR}^- \xrightarrow[\text{dry ether}]{\text{THF or}} \text{R}'\text{CH}=\text{CHR} + \text{Ph}_3\text{PO}$

50. Wohler's reaction

Acetylene is prepared by action of water on calcium carbide. This method is used for laboratory as well as industrial preparation.



51. Wurtz-Fittig Reaction

When a mixture of aryl halide and alkyl halide reacts with sodium metal in dry ether, substituted arenes are obtained.

